





### **CNMI Weekly Syndromic Surveillance Report**

**EPI WEEK DATE:** July 27, 2025 – August 02, 2025

Clinic	Influenza-Like-Illness (ILI)		Diarrhea (DIA)		Prolonged Fever (PF)		Acute Fever and Rash (AFR)		Total Encounters	
Cilife	Last week	Current week	Last week	Current week	Last week	Current week	Last week	Current week	Last week	Current week
CHCC Family Care Clinic	0	1	0	1	0	1	0	0	373	364
CHCC Women's Clinic	0	0	0	0	0	0	0	0	91	105
CHCC Children's Clinic	1	2	0	2	4	0	0	0	190	153
CHCC Emergency Room	10	9	6	6	8	6	0	0	338	331
Saipan Health Clinic	0	0	1	0	0	0	0	0	120	128
Kagman Isla Community Health	0	2	0	0	0	0	0	0	121	91
Southern Isla Community Health	Southern Isla Community Health 1		0	0	0	0	0	0	147	195
Tinian Isla Community Health	0	1	0	0	0	0	0	0	56	48
CHCC Lucia "Chiang" Villagomez Arizapa Health Center* 0		3	0	0	4	0	0	0	99	116
CHCC Rota Health Center	CHCC Rota Health Center 1 1		0	0	1	0	0	0	88	82
*CHCC Tinian Health Center was renamed to LCVA Health Center.	13	19	7	9	17	7	0	0	1623	1613

#### **ALERTS AND TRENDS**



ILI: Increase from previous week



**DIA: Stable** from previous week



**PF**: **Decrease** from previous week



AFR: Stable from previous week

#### **KEY TAKEAWAYS**

- **46% Decrease** in **Prolonged Fever cases** were seen this Epi Week (#31) compared to the average of the previous 3 Epi Weeks (#30, 29, & 28).
- > 36% Increase in Influenza-Like-Illness cases were seen this Epi Week (#31) compared to the average of the previous 3 Epi Weeks (#30, 29, & 28).
- > 35% Increase in Diarrhea cases were seen this Epi Week (#31) compared to the average of the previous 3 Epi Weeks (#30, 29, 28)

#### 1 Influenza case:

#### ❖ 1 Flu A H1-2009

Epi Week					Percent (%) change from	Antimicrobial Resistant (AMR) Infections				
Syndromes	31	30	29	28	current week to previous 3 weeks	Organism	EW 31	2025 YTD Totals		
Influenza-Like Illness	19	13	9	20	36%	MRSA	0	30		
Diarrhea	9	7	7	6	35%	VRE	0	3		
Prolonged Fever	7	17	14	8	-46%	ESBL	3	75		
Acute Fever and Rash	0	0	0	0	0%	CRE	0	0		







Commonwealth of the Northern Mariana Islands

### **CNMI Weekly Syndromic Surveillance Trends**

**EPI WEEK DATE:** July 27, 2025 – August 02, 2025











Commonwealth of the Northern Mariana Islands

### **CNMI Weekly Syndromic Surveillance Trends**

**EPI WEEK DATE:** July 27, 2025 – August 02, 2025









Commonwealth of the Northern Mariana Islands

### **CNMI Weekly Notifiable Disease Report for Select NNDs**

**EPI WEEK DATE:** July 27, 2025 – August 02, 2025

In the table below, weekly and year to date counts are displayed for Epi Week 31 and 2025, respectively. Additionally, a 3-year weekly average is calculated for the same Epi Week from the previous 3 years (2022-2024) for comparison to the current week. Incidence rates are calculated for 2024 and 2025 using the estimated population for the CNMI from the U.S. Census Bureau.

Condition	Epi Week 31	2025 YTD	3-year weekly average counts	2025 YTD Incidence Rates*	2024 Incidence Rates*
Enteric Diseases:					
Campylobacter	1	13	2	25.5	35.2
Ciguatera fish poisoning	0	5	0	9.8	9.8
Salmonella	0	13	0	25.5	43.0
Environmental:					
Elevated Blood Lead Levels	0	0	0	0.0	7.8
Sexually Transmitted Infections:					
Chlamydia	3	124	4	243.4	418.6
Gonorrhea	0	16	0	31.4	48.9
Syphilis	0	3	0	5.9	5.9
Respiratory Infections:					
Influenza	1	268	-	526.0	831.4
RSV	0	3	-	5.9	142.8
COVID-19	8	199	60	390.6	1299.0
Tuberculosis:					
TB, Confirmed	0	10	0	19.6	19.6
TB, Under Investigation	1	5	0	9.8	7.8

<sup>\*</sup>Rate per 100,000; Data are preliminary and subject to change. CNMI population estimates were determined using 2024 and 2025 Census International Database (<a href="https://www.census.gov/data-tools/demo/idb/#/country?YR">https://www.census.gov/data-tools/demo/idb/#/country?YR</a> ANIM=2021&COUNTRY YR ANIM=2021&FIPS SINGLE=CQ)








### **CNMI Weekly Prescription Drug Monitoring Program (PDMP) Report**

EPI WEEK 31 | EPI WEEK DATE: JULY 27 - AUGUST 2, 2025

WEEKLY CASE COUNTS											
POLYSUBSTANCE OPIOID			STIMULANT			BENZODIAZEPINE			OTHER SUBSTANCE		
OVERDOSE	MISUSE	OVERDOSE	OUD	MISUSE	OVERDOSE	StUD	MISUSE	OVERDOSE	BUD	MISUSE	OVERDOSE
0	2	0	0	0	0	0	3	0	0	0	1

NOTE: The encounters have been monitored since 2020. Some individuals might be involved in multiple cases or flagged multiple times for the same type of encounter in a single EPI week. The overdose Surveillance has expanded to include Stimulant and Polysubstance cases in 2021, Benzodiazepine cases in 2022. The Polysubstance cases are also counted under respective categories. Prior cases of any overdose related encounters might be duplicated under Other Substance Overdose category. Other Substance category analysis is solely depending on indications from the providers' notes. The substances reported are not verified by NDC number or DEA substance database.



OVERDOSE	Injury to the body (poisoning) that happens when a drug is taken in excessive amounts. An overdose can be fatal or nonfatal, accidental or intentional.							
	The use of more than one drug, also known as polysubstance use, is common. This includes when two or more are taken together or within a short time period, either intentionally or unintentionally. Intentional polysubstance use occurs when a person takes a drug to increase or decrease the effects of a							
POLY-SUBSTANCE	different drug or wants to experience the effects of the combination. Unintentional polysubstance use occurs when a person takes drugs that have been							
POLT-SUBSTANCE	mixed or cut with other substances, like fentanyl, without their knowledge. Whether intentional or not, mixing drugs is never safe because the effects							
	from combining drugs may be stronger and more unpredictable than one drug alone, and even deadly.							
	*For overdose Surveillance, Poly-Substance Use only includes encounters associated with Opioids, Stimulants, and/or Benzodiazepines.							
MISUSE	The use of illegal drugs and/or the use of prescription drugs in a manner other than as directed by a doctor, such as use in greater amounts, more often,							
IVIISUSE	or longer than told to take a drug or using someone else's prescription.							
OPIOID USE DISORDER	A problematic pattern of opioid, stimulant, or benzodiazepine uses that lead to serious impairment or distress. Diagnosing OUD/SUD/BUD requires a							
STIMULANT USE DISORDER	thorough evaluation, which may include obtaining the results of urine drug testing and prescription drug monitoring program (PDMP) reports, when							
DENIZODIAZEDINE LICE DICORDER	OUD/SUD/BUD is suspected. A diagnosis is based on specific criteria such as unsuccessful efforts to cut down or control use, or use resulting in social							
BENZODIAZEPINE USE DISORDER	problems and a failure to fulfill obligations at work, school, or home, among other criteria.							
	Any encounters that possibly leading to the above descriptions with such providers' comments as "requesting prescription refills (at emergency							
SUSPECTED MISUSE	department)", "drug-seeking-behavior", and "frequent ER visitor for the same complaint for chronic pain and requesting 'stronger' medication". Also,							
SUSPECTED MISUSE	cases where providers indicate there is possibility for misuse on the EHR system or when patients inform that they took Oxycodone (for example) and no							

PDMP data to support the patients' statement.

#### SENTINEL SITES

#### Commonwealth Healthcare Corporation (CHCC)

ER - Emergency Room, PCAP - Primary Care Access Point, CC - Children's Clinic, FCC - Family Care Clinic, WC - Women's Clinic,

THC - Tinian Health Clinic, RHC - Rota Health Center

Private Clinic

KICH - Kagman Isla Community Health, TICH - Tinian Isla Community Health, SICH – Southern Isla Community Health SHC – Saipan Health Clinic







### **CNMI Weekly Health & Vital Statistics Report**

### REPORTING PERIOD: EPI YEAR 2025 as of EPI WEEK 31

The statistics on births, deaths, and causes of deaths in this report are derived from birth and death registrations processed daily at the Health and Vital Statistics Office.

• Number	of births:	<b>8</b> <u>(314)</u>		•	Numbe	r of deaths	s:	<b>1</b> (1	<u>.36)</u>		
<ul> <li>Average</li> </ul>	Average: 10(per week)			•	Average: 4(per week)						
• Infection	Infections present and/or treated during					Number of deaths who received COVID-19 vacc					
pregnan	ıcy:										
o Chla	amydia:		<b>0</b> (8)		Age ran	ge:	< 5	≥ 5	12-17	18 & over	
o Gor	norrhea:		0(1)		N∘ of o	death	<b>0</b> (5)	<b>O</b> (0)	<b>O</b> (1)	<b>1</b> (130)	
o Syp	hilis:		<b>O</b> (0)		N∘ Va	ccinated	<b>O</b> (0)	<b>O</b> (0)	<b>O</b> (0)	<b>0</b> (100)	
o Hep	oatitis B:		0(1)		% Vac	cinated	0%	0%	0%	77%	
o Hep	oatitis C:		<b>1</b> (1)								
o CO\	/ID-19:		<b>O</b> (0)	•	<u>Mortalit</u>	ty Surveilla	ance:			<b>1</b> (136)	
Substan	ce use during	g pregnancy:			o Non-co	mmunical	ole disea	ses:		<b>1</b> (82)	
	rette smokin	· · ·	0(4)		•	Cancer rel	ated dea	ıths		<b>0</b> (20)	
o Bete	elnut chewing	g:	<b>0</b> (22)		•	Tobacco re	elated de	eaths		<b>0</b> (14)	
o Bete	elnut chewing	g + tobacco:	<b>0</b> (19)		o COVID-	-19 related	l deaths:			0(1)	
o Alco	o Alcohol use: <b>0</b> (1)					<b>O</b> (0)					
o Drug	-	abis, Crystal meth-	<b>0</b> (5)		_	COVID-13	other to	IIIIIDULIII	g conditions <sup>1</sup>		
	Ice, O	pioid, Others, etc.)			o Fetal D	eaths²:				<b>0</b> (5)	
o E-Ci	garette use:		<b>0</b> (2)								
	3 months b	efore pregnancy	<b>O</b> (0)		o Infant I	Deaths:				<b>0</b> (5)	
•	During preg	gnancy	<b>0</b> (2)		o Childre	n (aged 1	- 4 years	) Deaths	:	<b>o</b> (0)	
		in pregnancy:			o Materr	nal Deaths:		•		<b>0</b> (0)	
	pregnancy D	M:	<b>0</b> (5)								
	tational DM:		<b>1</b> (42)			nt or Injur	-	d Deaths	.3. ) .	<b>0</b> (5)	
	pregnancy H		0(11)			Drowning:				0(1)	
	tational HTN:	•	<b>1</b> (25)			Suicide:				0(4)	
1	•	ow survival births)				Homicide:				<b>0</b> (0)	
	h weight < 15	-	<b>1</b> (3)		•	Traffic fata	ality:			<b>0</b> (0)	
1	h weight < 25	•	<b>2</b> (25)		•	Drug and/	or opioid	doverdo	se:	0(1)	
o Ges	tation age < 3	37 weeks:	<b>1</b> (31)		•	Poisoning:	:			<b>O</b> (0)	

¹ Other significant condition contributing to death but NOT resulting in the underlying cause.² Fetal deaths = Fetus weighed ≥ 350 grams, or fetal demise > 20 weeks of completed gestation.

Data source: Electronic Vital Registration System (EVRS)

<sup>&</sup>lt;sup>3</sup> Accident or Injury related deaths = Manner of death beyond Natural Causes: accidental deaths (traffic and drowning), suicide, drug overdose, and poisoning.








### **CNMI Weekly Health & Vital Statistics Report**

#### REPORTING PERIOD: EPI YEAR 2025 as of EPI WEEK 31

The statistics on births, deaths, and causes of deaths in this report are derived from birth and death registrations processed daily at the Health and Vital Statistics Office.



<sup>4</sup>Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified; <sup>5</sup> Mental, Behavioral and Neurodevelopmental disorders; <sup>6</sup>Certain conditions originating in the perinatal period; <sup>7</sup>Congenital malformations, deformations and chromosomal abnormalities; <sup>8</sup>Injury, poisoning and certain other consequences of external causes; <sup>9</sup>Diseases of the musculoskeletal system and connective tissue, <sup>10</sup>Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism

Vital events reported, December 29, 2024 - August 02, 2025

